



Specimen Label

- May be used on Livestock Animals and Livestock Premises
- May be used in Food Handling Establishments
- Kills Bed Bugs
- Kills Bees and Wasps [Yellow Jackets, Hornets and Paper Wasps]

ACTIVE INGREDIENTS:

*Pyrethrins.....	1.00%
OTHER INGREDIENTS	99.00%
	100.00%

*Pyrethrins are a botanical insecticide derived from chrysanthemum flowers.

0.01 lbs. active ingredient per lbs. of Evergreen® Pyrethrum Dust



KEEP OUT OF REACH OF CHILDREN
CAUTION

See inside for first aid and precautionary statements.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL BEFORE EACH USE

USE RESTRICTIONS:

- Apply this product only as specified on the label.
- Aerial applications are prohibited.
- Do not allow contamination of food and feedstuffs.
- In indoor residential areas or other areas where children may come into contact, only apply as a crack-and-crevice or as a spot treatment to areas no greater than 3 feet by 3 feet per room.
- Applications with power duster equipment are prohibited.
- Do not enter or allow others to enter until dusts have settled.
- Do not apply this product in a way that will contact workers or other persons, either directly or indirectly through drift.
- Only protected handlers may enter treated area during application.
- Do not allow adults, children or pets to enter the treated area until dusts have settled.
- For use outdoors, do not apply directly to or near water, storm drains or drainage ditches. Do not apply when windy.
- Do not water the treated area to the point of runoff.
- Do not make applications during rain.
- Do not apply this product in a way that will contact adults, children or pets, either directly or indirectly through drift.
- Remove or cover exposed food and drinking water before application.
- Remove or cover dishes, utensils, food processing equipment and food preparation surfaces, or wash them before use. Do not use in serving areas while food is exposed.

- In Food Areas of Food Handling Establishments: Do not contaminate food or feedstuffs. During applications, cover or remove food. Cover food processing surfaces and equipment or clean after treatment with a suitable detergent and rinse with potable water before use. Do not use in serving areas (dining areas where prepared foods are served) while food is exposed or facility is in operation.
- In Schools, do not apply to classrooms when in use.
- Do not use in aircraft cabins.
- Remove pets, birds and cover fish aquariums before application.
- When used in dairy barns or facilities; close milk bulk tank lids to prevent contamination from spray and from dead or falling insects. Remove or cover milking utensils before application. Wash teats of animals before milking.
- Except when applying to sewers or drains as instructed on this label, do not apply directly into sewers or drains, or to any area like a gutter where drainage to storm sewers, water bodies, or aquatic habitat can occur. Do not allow the product any drain during or after application.
- Not for use on commercial sod farms.
- **USE RESTRICTIONS IN FOOD HANDLING AREAS:**
- Do not apply when food or feed processing facility is in operation or foods are exposed.
- **APPLICATIONS OF THIS PRODUCT IN FOOD/FEED AREAS OF FOOD/FEED HANDLING ESTABLISHMENTS ARE NOT PERMITTED, OTHER THAN AS A CRACK-AND-CREVICE TREATMENT.**

Phytotoxicity:

Plant safety is an important consideration when using insecticides. It is not possible to evaluate the phytotoxicity of Evergreen® Pyrethrum Dust on numerous plant varieties that may react differently to insecticides in different growth stages or under varying environmental conditions. Before making widespread applications of Evergreen® Pyrethrum Dust treat a limited number of plants and observe for phytotoxicity over a 10-day period.

Greenhouses:

This product can be used in greenhouses to kill listed insects. Apply to cover upper and lower surfaces of leaves stems and flowers with a thin even film of dust. Apply Evergreen® Pyrethrum Dust to ornamental plants or only to fruit, vegetable and herb crops in the plug stage prior to transplant. Do not apply to fruit, vegetable, or herb crops that are ready for consumption.

OUTDOOR AREAS for use in and around areas:

Ant Mounds, Arboretums, Backyards, Balconies, Beneath Decks, Beneath Gazebos, Calf Hutches, Compost Sites, Courtyards, Decks, Dog Runs, Drive-In Restaurants, Drive-In Theaters, Dumpsters, Farmers Markets, Flea Markets, Front Yards, Gardens, Gazebos, Home Gardens, Kennels, Lanais, Landscapes, Lath Houses, Loading Docks, Lumber Storage Areas, Nurseries, Outdoor Eating Facilities, Patios, Porches, Sanitary Sewers (Manholes), Side Yards, Swine Yards and Wood Piles.

OUTDOOR APPLICATIONS	
Application Type	Application Rate
Asian Lady Beetles and Box Elder Bugs – Perimeter Building Exterior	16 to 24 oz. / 1,000 sq. ft.
Ornamental Plants	Apply to cover upper and lower surfaces of leaves, stems and flowers, with a thin even film of dust.
Insects Harboring in Lawns, Landscaping and Backyards	Apply 8 to 24 oz. / 1,000 sq. ft.
Building Perimeter Treatments	Apply 16 to 24 oz. / 1,000 sq. ft.

Building Perimeter Treatments:

To aid in preventing entry and infestations of buildings by listed insects, apply Evergreen® Pyrethrum Dust to building exterior as a perimeter treatment in a band extending 12"-18" along and adjacent to foundation, and to a height of 2 to 3 feet up the side of the building where these pests are active and may find entrance.

To Kill Insects Harboring in Lawns, Landscaping, and Backyards:

To aid in the kill of listed insects, apply Evergreen® Pyrethrum Dust at the specified rate. For best results, mow lawn one to two (1-2) days before applying, and remove clippings. Be sure to cover the entire lawn surface where insects are present, or normally frequent. Apply around the base of ornamental plants, shrubs and trees. Do not water or irrigate immediately following application. Keep pets and children out of the treated area until after the dust has settled.

LIVESTOCK, DAIRY, SWINE, POULTRY AND OTHER ANIMAL BUILDINGS AND PREMISES

LIVESTOCK PREMISES	
Application Type	Application Rate
Crack and Crevice	1 oz. / 50 sq. ft.
Spot Treatments and Surface Treatments	16 to 24 oz. / 1,000 sq. ft.
Building Perimeter Treatments	16 to 24 oz. / 1,000 sq. ft.
Spot Treatments for Ticks	16 to 24 oz. / 1,000 sq. ft.

Kills: Bed Bugs, Blacklegged Tick Nymphs, Chicken Mites, Deer Ticks, Face Flies, Fleas, Flies, Northern Fowl Mites, Poultry Lice and Stable Flies

To aid in the kill of listed insects in listed indoor and outdoor animal areas, apply Evergreen® Pyrethrum Dust to cracks, crevices, voids, structural joints, floors, walls, wall joints, support beams, trusses, rafters, ceilings, around scales, chutes, gates, and other areas where insects harbor, alight, hide or travel. For Ticks, apply Evergreen® Pyrethrum Dust in areas where they like to harbor such as cracks and crevices.

Apply Evergreen® Pyrethrum Dust to exterior of livestock buildings in an 18" band along and adjacent to building foundations, and to a height of 2 to 3 feet up the side of the building where insects are active and may find entrance.

LIVESTOCK ANIMAL USE

LIVESTOCK ANIMALS	
Application Type	Application Rate
Beef, Dairy Cattle, Horses	2 oz. dust per animal
Swine	1 oz. per head as a uniform coat to head, shoulders and back.
Poultry	1 lb. / 100 birds.

Kills: Bed Bugs, Blacklegged Tick Nymphs, Chicken Mites, Deer Ticks, Face Flies, Fleas, Flies, Northern Fowl Mites, Poultry Lice and Stable Flies

Beef, Dairy, Cattle, Horses and other Livestock:

Can be used in dust bags, shaker can and mechanical dust applicator to aid in the kill or repellency of listed livestock insects.

Direct Application: Apply 2 oz. of dust per animal by shaker can, or suitable mechanical dust applicator over the head, neck, shoulders, back and legs.

Application through Dust Bags: Place contents of this package in any commercially available dust bag, suspend bag in areas frequented by cattle or in gateways or lanes through which the animals must pass daily for water, feed or minerals. Bags may also be placed in loafing sheds or in front of mineral feeders. For dairy cows, bags may be suspended in the exit through which the cows leave the milking barn. Hang the bags 4 to 6 inches below the back line of the cattle. For reduction of Face Flies, bags must be located so animals will be forced to use them daily and hung at a height so that the face is dusted.

Swine:

USE RESTRICTIONS FOR APPLICATION TO SWINE

- Do not apply more often than once every 10 days
- Do not ship animals for slaughter within 5 days of treatment.

Direct Application: Apply only 1 oz. per head as a uniform coat to the head, shoulder and back by use of a shaker can or suitable mechanical dust applicator. In severe infestation, both animals and the bedding may be treated.

Poultry:

To kill listed livestock insects, apply at a rate of 1 lb. per 100 birds.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE:

Keep this product in its tightly closed original container, when not in use. Do not store near heat or open flame. Store in a cool, dry, well-ventilated (preferably locked) area that is inaccessible to children and animals, and away from food and pet food.

PESTICIDE DISPOSAL:

To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

CONTAINER DISPOSAL:

Nonrefillable container. Do not reuse or refill this container. Completely empty container by shaking and tapping sides and bottom to loosen clinging particles promptly after emptying. Then offer for recycling if available or reconditioning, if appropriate or dispose of in a sanitary landfill or by other procedures approved by State and Local authorities.

DIRECTIONS FOR USE CONTINUED

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval of 12 hours.

PPE required for early entry to treated areas (that is permitted under Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water), is:

- Long-sleeved shirt and long pants;
- Shoes plus socks;
- chemical-resistant gloves made of any waterproof material.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR, Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow others to enter until dusts have settled.

Kills Exposed Listed Insects:

Adelgids	Brown Spider Beetles	Crickets	Flying Moths	Larger Cabinet Beetles	Pine Engraver	Southern Pine Beetles
Alfalfa Weevil	Brown Spruce Longhorn Beetle	Cross-striped Cabbageworm	Foreign Grain Beetles	Beetles	Pink Scavenger Caterpillars	Southern Red Mite (Adult)
Almond Moth	Budworms	Cucumber Beetles	Forest Tent Caterpillar	Larger Grain Borers	Pinworms	Sowbugs
American Black Flour Beetle	Cadellles	Cutworms (exposed)	Fruit Flies	Leafhoppers	Plant Bugs	Spider Beetles
Angoumois Grain Moths	California Oakworm	Dark Mealworms	Fruittree Leafroller	Leafminers	Plum Curculio	Spider Mites
Ants	Cankerworms	Darkling Beetles (lesser meal worm)	Fungus Beetles	Leafrollers	Ponderosa Pine Needle Miner	Spruce Mite (Adult)
Aphids	Carpet Beetles	Deer Tick	Fungus Gnats	Leaf tiers	Poultry Lice	Squarenecked Grain Beetles
Apple Maggot	Carrion Beetles	Depressed Flour Beetles	German Cockroaches	Lesser Grain Beetles	Powder Post Beetles	Stable Flies
Armyworms	Carrot Weevil	Dermestid Beetles	Giant Root Aphid	Lesser Grain Borers	Psocids	Stalk Borers
Asian Cerambycid Beetles	Caterpillars	Diamondback Larvae	Glassy Winged Sharpshooter	Lesser Meal Worm	Psyllids	Stink Bugs
Asian Lady Beetle	Catorama Beetles	Dingy Cutworm Adult	Grain Mites	Lesser Peach Tree Borer	Red Flour Beetles	Stored Product Pests
Asian Pine Tip Moths	Centipedes	Douglas Fir Midge	Granary Insects	Longheaded Flour Beetles	Red Gum Lerp Psyllid	Subtropical Pine Tip Moth
Bagworm	Cereal Beetles	Douglas Fir Tussock Moth	Granary Weevils	Loopers	Red Horned Grain Beetles	Sycamore Leaf Beetle
Barn Flies	Cheese Mites	Drain Flies	Grape Leaf Skeletonizer	Lygus Bugs	Red Lily Beetles	Tarnished Plant Bug
Bat Bugs	Cheese Skippers	Dried Fruit Beetles	Grape Leafhopper	Maize Weevils	Red Oak Borer	Tent Caterpillar
Bean Beetles	Chicken Mites	Drugstore Beetles	Grasshoppers	Mealy Bugs	Redbanded Leafroller	Thrips
Bean Weevils	Chrysanthemum Leaf Miners	Earwigs	Green Fruit Worm	Mediterranean Flour Moths	Rice Flour Beetles	Tobacco Moths
Bed Bugs	Cicadas	Eastern Tent Caterpillar	Green Peach Aphid	Merchant Grain Beetles	Rice Moths	Trogoderma Beetles
Bees	Cigarette Beetle	Elmleaf Beetle	Greenbugs	Mexican Bean Beetle	Rice Weevils	Two-Banded Beetles
Beet Armyworms	Clearwing Borer	Emerald Ash Borer	Greenhouse Thrips	Mexican Grain Beetles	Root Weevils	Vegetable Leaf Miners
Beetles	Clothes Moths	European Grain Moths	Ground Beetles	Millers	Rose Chafer	Vinegar Flies
Birch Leafminer	Clover Mites	European Pine Shoot Moth	Gypsy Moth (adults & larvae)	Millipedes	Royal Palm Bugs	Warehouse Beetles
Black Carpet Beetles	Clover Weevil	European Pine Tip Moth	Hairy Fungus Beetles	Mimosa Webworm	Rusty Grain Beetles	Wasps [Yellow Jackets, Hornets and Paper Wasps]
Black Fungus Beetles	Cluster Flies	European Pine Tip Moth	Hairy Spider Beetles	Mole Crickets	Saw Flies	Waterbugs
Black Turpentine Beetles	Cocoa Bean Moths	Face Flies	Harlequin Bug	Moth Flies	Saw-Toothed Grain Beetle	Webworms
Black Vine Weevil	Coffee Bean Weevils	Fall Cankerworms	Heliolithis spp.	Murmidius Beetles	Scale (Crawlers)	Weevils
Blacklegged Tick Nymphs	Cone Beetles	Fall Webworm	Hessian Fly (Adult)	Mushroom Flies	Seedbugs	Western Pine Tip Moth
Blister Beetles	Cone Worms	Fannia Flies	Hide Beetles	Nantucket Pine Tip Moth	Silverfish	White Apple Leafroller
Blow Flies	Confused Flour Beetle	Field Crickets	Hornworm	Northern Fowl Mites	Skipper Flies	White Marked Spider Beetles
Booklice	Conifer Spider Mite (Adult)	Firebrats	Inchworm	Onion Maggot	Skippers	Whiteflies
Bottle Flies	Corn Sap Beetles	Fireworms	Indian Meal Moth	Palmetto Bug	Slenderhorned Flour Beetles	Wireworms and Yellow Mealworms
Box Elder Bugs	Cottonwood Borer	Flat Grain Beetles	Iris Borer	Peach Twig Borer	Small Flying Moths	
Broadhorned Flour Beetles	Cottonwood Leaf Beetles	Flea Beetles	Japanese Beetle	Pear Psylla	Small Fruit Flies	
Broadnosed Grain Weevils	Cranberry Girdlers	Fleas	Katydids	Pillbugs	Smalleyed Flour Beetles	
	Crane Flies	Flies	Khapra Beetles	Pine Bark Beetles		
			Lace Bugs			
			Large Black Flour Beetles			

Application Methods:

Apply Evergreen® Pyrethrum Dust using a non-power dust applicator, a shaker can, dust gun, puffer or other bulbous duster, hand pump duster, hand-operated rotary duster, static charge duster, or other appropriate hand-operated duster.

INDOOR AREAS: For use in and around indoor areas:

Animal Areas	Coffee Houses	Egg Processing Plants	Goat Houses	Livestock Hauling Equipment	Pet Food Plants	Swine Houses
Animal Quarters	Coffee Shops	Egg Storage Facilities	Grain Elevators	Livestock Holding Areas	Pet Sleeping Areas	Taverns
Apartments	Commercial Buildings	Factories	Grain Handling Equipment	Poultry Houses	Poultry Processing Plants	Tea Houses
Asian Markets	Condominiums	False Ceilings	Grain Harvesting Equipment	Rabbit Houses	Rabbit Processing Plants	Textile Mills
Attics	Convenience Stores	Farm Structures	Grain Mills	Railroad Cars	Research Animal Areas	Tobacco Factories
Bakeries	Crawl Spaces	Feed Mills	Grain Storage	Restaurants	Rice Mills	Tobacco Warehouses
Barns	Dairies	Food Handling Establishments	Granaries	Schools	Seed Storage	Tool Sheds
Basements	Dairy and Livestock Barns	Food Preparation Facilities	Greenhouses	Shade Houses	Ships	Trains
Beef Barns	Dairy Food Processing	Food Processing Facilities	Grocery Stores	Ships Holding Areas	Ships	Transportation Equipment (except Aircraft)
Beverage Plants	Day Care Centers	Food Processing Facilities	Hog Barns	Ships	Ships	Tree Nuts Storage Areas
Bins	Delicatessens	Food Quarantine Areas	Homes	Shops	Shops	Truck Trailers
Bottling Plants	Department Stores	Food Service Facilities	Horse Stables	Shops	Shops	Vegetable Packing Facilities
Brandy Storage Warehouses	Diners	Food Storage Facilities	Horse Stables and Barns	Shops	Shops	Vegetable Processing Facilities
Breweries	Distilleries	Fruit Packing Sheds	Horse Trailers	Shops	Shops	Vegetable Storage Facilities
Brewery Warehouses	Dog Houses	Fruit Processing Plants	Hospitals	Shops	Shops	Veterinary Clinics
Buildings Housing Animals	Dog Sleeping Quarters	Fruit Quarantine Areas	Hotels	Shops	Shops	Veterinary Hospitals
Buildings Housing Poultry	Drains & Sewers	Fruit Storage Warehouses	Houses	Shops	Shops	Voids in Attics
Cabinets	Dried Flower Warehouses	Garbage Bins	Indoor Eating Facilities	Shops	Shops	Wall Voids
Cafeterias	Dried Food Products in Processing	Garbage Cans	Industrial Installations	Shops	Shops	Warehouses
Calving Barns	Dried Food Products in Storage	Garbage Compactors	Industrial Plants	Shops	Shops	Wheat Mills
Canneries	Dried Fruit Processing Plants	Garbage Trucks	Institution Dining Areas	Shops	Shops	Wholesale Stores
Cat Sleeping Quarters	Dried Fruit Warehouses	Goat Holding Areas	Institutions	Shops	Shops	Wine Cellars
Cattle Barns	Dumpsters		Kennels	Shops	Shops	Wine Storage Warehouses
Certified Organic Facilities	Dwellings		Liveries	Shops	Shops	Wine Tasting Rooms
Cheese Caves			Livestock Barns	Shops	Shops	Wineries and Zoos
Citrus Packing Houses				Shops	Shops	

APPLICATIONS WITHIN FOOD HANDLING ESTABLISHMENTS

Food and/or Feed areas include areas for receiving, storage (dry, cold, frozen or raw), packaging (canning, bottling, wrapping, boxing and grinding), preparing, edible waste storage and enclosed processing systems (mills, dairies, and systems for processing edible oils and syrups). Serving areas are places where prepared foods are served.

FOOD HANDLING ESTABLISHMENTS	
Application Type	Application Rate
Crack and Crevice Food/Feed Areas of Food/Feed Handling Establishments	1.5 oz. / 50 sq. ft.
Crack and Crevice Stored Product Pests in Non-Food/ Non-Feed Areas of Food Handling Establishments	1.0 oz. / 50 sq. ft.
Crack and Crevice Treatments in Storage Sites	1.0 oz/ 50 sq. ft.

Using a non-power duster, apply directly into cracks and crevices, in expansion joints between different elements of construction, under or between equipment bases and the floor, beneath appliances, into wall voids, motor housing, junction boxes or switch boxes, conduits, behind and under cabinets or drawers, in and around drains, around drain pipes, plumbing and utility entrances, hollow equipment legs and other sites where listed insects can enter or hide.

Stored Product Pests in Non-Food/Non-Feed areas of Food Handling Establishments:

Non-food/feed areas of food/feed handling establishments are garbage rooms, lavatories, floor drains (to sewers, entries and vestibules), offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after canning or bottling).

To aid in the kill of stored product pests in listed areas. Apply Evergreen® Pyrethrum Dust at the specified rate. Clean up waste materials, infested materials, dust, dirt, and any other debris. Thoroughly treat floors, walls, shelves, drawers, cabinets and other surfaces in bins, storage and handling areas, around cartons or containers and other areas where these insects tend to congregate. Apply into cracks and crevices and other difficult-to-reach areas where insects may hide. Apply in voids or cracks and crevices in pantries and other infested areas within the structure.

For Use in Storage Sites:

This product can be used in warehouse bins, trucks, cargo ships, mills, bin hoppers, elevators and conveying equipment, and other listed sites as a cleanup prior to using them for storage. Remove and dispose of all of the infested stored commodity accumulations and debris, with special attention to material lodged in the cracks and crevices. After sanitation measures have been employed, dust areas prior to use for storage, paying particular attention to cracks and crevices.

For packaged food warehouses, treat potential points of insect entry, as well as harborage sites in, on and around wooden and plastic pallets, pallet-racking systems, loading docks, scales, around cartons.

INDOOR TREATMENTS

IN INDOOR RESIDENTIAL AREAS OR OTHER AREAS WHERE CHILDREN MAY COME INTO CONTACT, ONLY APPLY AS A CRACK-AND-CREVICE OR AS A SPOT TREATMENT TO AREAS NO GREATER THAN 3 FEET BY 3 FEET PER ROOM.

APPLICATION RATES FOR INDOOR TREATMENTS		
Application Type	Application rates for Residential areas such as homes, condominiums, apartments, hotels, motels and pet areas	Application rates for Non-Residential areas such as offices, warehouses, hospitals and other commercial buildings and institutions
Crawling Insects—Crack and Crevice	Apply as a crack-and-crevice treatment to areas no greater than 3 x 3 feet per room.	1.5 oz. / 50 sq. ft.
Bed Bugs—Crack and Crevice	Apply as a crack-and-crevice treatment to areas no greater than 3 x 3 feet per room.	2.0 oz. / 50 sq. ft.
Ants—Crack and Crevice	Apply as a crack and crevice treatment to areas no greater than 3 x 3 feet per room.	1.0 oz. / 50 sq. ft.
Crawling Insects for Severe Infestations—Spot Treatments	Apply as a crack and crevice or as a spot treatment to areas no greater than 3 x 3 feet per room.	16 to 20 oz. / 1,000 sq. ft.
Fleas, Ticks and Mites in Areas where Pets Frequent—Spot Treatments	Apply as a spot treatment to areas no greater than 3 x 3 feet per room.	1.0 oz. / 70 sq. ft.
Powder Post Beetles —Spot Treatments	Apply as a spot treatment to areas no greater than 3 x 3 feet per room.	16 to 20 oz. / 1,000 sq. ft.
Asian Lady Bird Beetles and Box Elder Bugs in Crawl Spaces and Attics—Crack and Crevice	Apply as a spot treatment to areas no greater than 3 x 3 feet per room.	1 oz. / 50 sq. ft.
Asian Lady Bird Beetles and Box Elder Bugs in Crawl Spaces and Attics—Spot Treatments	Apply as a spot treatment to areas no greater than 3 x 3 feet per room.	16 to 24 oz. / 1,000 sq. ft.
Ornamental Plants	Apply to cover upper and lower surfaces of leaves, stems and flowers, with a thin even film of dust.	Apply to cover upper and lower surfaces of leaves, stems and flowers, with a thin even film of dust

Crawling Insects:

Apply Evergreen® Pyrethrum Dust into voids, cracks and crevices or other places where insects are traveling or harboring. Apply a continuous visible film underneath and behind stairs, behind and beneath sinks, stoves, refrigerators and other appliances, around garbage cans, cabinets. Treat on, around and behind moldings, shelving, baseboards, doorsills, doorframes, floors, around drains, pipes and pipe openings, plumbing, bookcases, storage, utility installations and other areas where insects hide or travel. For severe infestations, apply behind cabinets and walls, in crawl spaces and attics at the specified rate.

Fleas, Ticks, and Mites:

DO NOT TREAT DOGS OR CATS WITH THIS PRODUCT.

For Fleas, apply Evergreen® Pyrethrum Dust only as a direct treatment. Treat on and around animal bedding, sleeping quarters, kennel, furniture, floor and floor coverings and all areas where these pests inhabit. Treat cracks and crevices behind and around window and door frames, wall cracks, moldings, shelving, baseboards, pipe openings and all other cracks. Treat the entire inside surface of doghouses. For Ticks, apply Evergreen® Pyrethrum Dust in places where the insects are traveling or harboring.

Bed Bugs:

Thoroughly inspect area of suspected bed bug infestation, and note potential harborage sites or actual infestations. Remove and/or pick up clothing and other obstacles that may prevent thorough inspection or treatment. Remove and sanitize infested bedding (pillows and linens), clothing, or other items.

Take bed apart. Apply Evergreen® Pyrethrum Dust into joints and channels if hollow, such as square or round tubing, and into cracks and crevices on the interior of bed frame, headboards, and footboards. If hollow, ensure that the interior framework is well dusted. Wall-mounted headboards and their hardware should be removed and treated. Mattresses should be dusted, especially seams, edges, and tufts. Mattress treatments must be limited to seams, folds and edges only.

Bed bugs may also harbor in areas of the room away from the bed. Apply Evergreen® Pyrethrum Dust as a crack and crevice application where evidence of bed bugs occurs. All furniture and furniture accessories in the room must be treated, including inside empty drawers and shelving in clothes closets, night stands and other furniture in the room, carpet edges, high and low wall moldings (such as floor molding or window casings or ledges/sills), door jambs and hinges, wall voids, and behind wall hangings and mirrors. Pull carpeting from tack strip and treat under the baseboard, sill plate, and under the carpet tack strips. Remove all clothes and other articles from dressers or closets before application. Treat all cracks and crevices associated with light switches, wall plates, faceplates, outlet covers, mounting boxes and housings. Apply Evergreen® Pyrethrum Dust as a continuous film to other potential harborage sites, such as table and wall-mounted lamps, wall crevices, drapery rods, curtains, blinds/mounting hardware, and other cracks and crevices in the room.

Ants:

To aid in the kill of listed Ants, apply Evergreen® Pyrethrum Dust at the specified rate. Treat door frames, around window frames, behind, on and around moldings, cabinets, pipe openings, utility openings, shelves, ant trails, hills, wall voids, cracks and crevices and other points of entry into premises. Combine interior treatments with exterior perimeter application of Evergreen® Pyrethrum Dust.

Powder Post Beetles*:

To aid in protecting against insect entry. Apply to distribute Evergreen® Pyrethrum Dust through insect entry points, crawl holes into attics and crawl spaces of buildings. Close all vents and blowers so that all target surfaces can be treated. Contact all wood surfaces. In case of localized infestations, probe to locate galleries and drill holes to inject Evergreen® Pyrethrum Dust into the galleries and void areas. If a heavy infestation is suspected, a conventional, industry accepted techniques and treatments should be employed.

*The purpose of such applications of Evergreen® Pyrethrum Dust is to kill workers or winged reproductive forms that may be present in treated channels at the time of treatment. Such applications are not a substitute for mechanical alteration, soil treatment or foundation treatment, but are merely a supplement. This product is not for use against Termites. For active Termite infestations, use conventional, industry accepted techniques and treatments.

Asian Lady Beetles and Box Elder Bugs:

To kill or repel these insects, as well as Clover Mites, Cluster Flies and Elmleaf Beetles, inject Evergreen® Pyrethrum Dust into cracks and crevices and/or voids where these insects may be entering the premise, harboring or hibernating. Apply in crawl spaces and attics. On building exterior, treat underneath wood fascia, behind siding, cracks around windows, doors, siding, wall voids, utility pipes and entry points, behind chimneys, around foundations and foundation cracks, patios and deck joints. When treating for Asian Lady Beetles, apply Evergreen® Pyrethrum Dust in, on and around areas with abrupt color contrasts on buildings (shutters, drainpipes, etc), southwest-facing or sun-exposed sides of buildings, and vertical corners of buildings.

Ornamental Trees, Shrubs, Flowers and Foliage Plants: Including: African Violet, Ageratum, Andromeda, Arborvitae, Ash, Aster, Azalea, Beech, Begonia, Birch, Boxwood, Calceolaria, Calendula, Calla, Camellia, Carnation, Ceanothus, Chrysanthemum, Cineraria, Coleum, Cotoneaster, Crabapple, Cyclamen, Cypress, Daffodil, Dahlia, Delphinium, Dogwood, Elm, Eucalyptus, Euonymus, Fern, Ficus, Fir, Firethorn, Foliage Plants, Forsythia, Fuchsia, Gardenia, Geranium, Gladiolus, Gloxinia, Gypsophila, Hawthorn, Hemlock, Hickory, Holly, Honey Locust, Horse Chestnut, Hyacinth, Hydrangea, Iris, Juniper, Larch, Laurel, Lilac, Lily, Linden, Maidenhair Fern, Marigold, Mimosa, Myrtle, Narcissus, Oak, Palm, Pansy, Pelargonium, Peony, Petunia, Philodendron, Phlox, Pine, Privet, Pyracantha, Rhododendron, Roses, Rubber Plant, Snapdragon, Sweet Pea, Tulip Tree, Tulips, Viburnum, Wandering Jew, Willow, Yew, Zinnia.

Evergreen® Pyrethrum Dust must come into contact with insects in order to kill them. Coverage of the target insect is vital to achieving good performance. Apply Evergreen® Pyrethrum Dust to cover upper and lower surfaces of leaves stems and flowers with a thin, even film of dust. Apply when insects or insect damage first appears.

When treating trees, shrubs and similar plants, treat the trunk, the base of the trunk and the critical root zone. Treat the tree crotch, knotholes, wounds and other areas where insects may harbor. Allow dust to settle in treated areas before reentering. Do not water or irrigate immediately following treatment. Do not eat or consume treated plants for 1 year after treatment.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Mixers, Loaders, Applicators, and other handlers must wear the following:

- long-sleeved shirt
- long pants
- shoes and socks
- chemical-resistant gloves made of any waterproof material

USER SAFETY REQUIREMENTS

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Discard clothing and other absorbent material that have been drenched or heavily contaminated with the product's concentrate. Do not reuse them.

USER SAFETY RECOMMENDATIONS

Users should: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic organisms, including fish and invertebrates. To protect the environment, do not allow pesticide to enter or runoff into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area.

This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

KEEP OUT OF REACH OF CHILDREN

CAUTION PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.

(If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID

IF ON SKIN OR CLOTHING:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For additional information on this pesticide product (including health concerns, medical emergencies or pesticide incidents), you may call 1-888-740-8712.

